

Prevalence and Risk Factors of Apnea in Preterm Neonates Admitted to the French Medical Institute for Mothers and Children Hospital in Kabul City: An Analytic Cross-Sectional Study

Mansoor Aslamzai MD^{1*}, Basir Ahmad Froogh MD², Omid Faizi MD³, Mohammad Farouq Hamidi MD⁴ and Sayed Azam Sajid MD⁵

¹Professor; Department of Neonatology, Kabul University of Medical Science, 3rd district, 1003, Kabul, Afghanistan.

²Associate Professor; Department of Neonatology, Kabul University of Medical Science, 3rd district, 1003, Kabul, Afghanistan.

³Neonatologist and DCH, Chief of Neonatology Department, French Medical Institute for Mothers and Children Hospital, 3rd district, 1003, Kabul, Afghanistan.

⁴Associate Professor; Department of Neonatology, Kabul University of Medical Science, 3rd district, 1003, Kabul, Afghanistan.

⁵Clinical Documentation Specialist, Saints John's Regional Medical Centre, United States.

*Correspondence:

Mansoor Aslamzai, MD, DCH Trainer, Professor of Neonatology, Kabul University of Medical Science, 3rd district, 1003, Kabul, Afghanistan. Email: mansoraslamzai@gmail.com. Tel: 0093700603998- 093788449693.

Received: 16 May 2022; Accepted: 22 Jun 2022; Published: 27 Jun 2022

Citation: Aslamzai M, Froogh BA, Faizi O, et al. Prevalence and Risk Factors of Apnea in Preterm Neonates Admitted to the French Medical Institute for Mothers and Children Hospital in Kabul City: An Analytic Cross-Sectional Study. J Pediatr Neonatal. 2022; 4(2): 1-5.

ABSTRACT

Introduction: Prematurity is one of the leading causes of neonatal death in Afghanistan and complicates many serious problems including apnea. Although recurrent and prolonged apnea of prematurity (AOP) may be directly or indirectly associated with significant adverse outcomes, there isn't enough scientific information regarding the prevalence and risk factors of this problem in Afghanistan, therefore, this study is performed to obtain such information.

Objective: To determine the prevalence rate and risk factors for apnea of prematurity in neonates admitted to the Neonatal Intensive Care Unit of the French Medical Institute for Mothers and Children Hospital, Kabul City.

Patients and Methods: This analytic cross-sectional study was conducted at the Neonatal Intensive Care Unit of the French Medical Institute for Mothers and Children Hospital in Kabul City, Afghanistan. The study participants were preterm neonates. Statistical analysis was performed by SPSS 24.

Results: 75 preterm newborns were enrolled in this study and the apnea of prematurity were developed in 48% of them. Extremely low birth weight preterm neonates had a 71.4% prevalence rate of such apnea, while in very low birth weight and low birth weight the rates were 47.4% and 36.4% respectively. Based on gestational age, the prevalence rates of apnea in the early and moderate preterm neonates were 55.6% and 52.8% respectively, whilst also in late preterm neonates it was 42.8%. The apnea of prematurity appeared more prevalent in boys (54.3%) than girls (45.7%). The preterm neonates in the apnea group versus the non-apnea group had the mean birth weight of (1233.33 ± 235.25g vs 1333.46 ± 274.44g, 90%CI= -198 _ -1.4), mean maternal age of (24.78 ± 3.68y vs 26.62 ± 4.58y, 90%CI= -3.44 _ -0.23) and RR of anemia (2.2, P=0.05).

Conclusion: The overall prevalence rate of AOP in preterm neonates was 48% and the highest rates were seen within extremely low birth weight and early preterm neonates. The lower neonatal birth weight, neonatal anemia and younger maternal age were found to be the risk factors for apnea of prematurity.

Keywords

Cessation of Respiration, Newborn, Prematurity.

Abbreviations

AOP: Apnea of Prematurity, ELBW: Extremely Low Birth Weight, FMIC: French Medical Institute for Children and Mother, IVH: Intraventricular Hemorrhage, LBW: Low Birth Weight, LMP: Last Menstrual Period, NEC: Necrotizing Enterocolitis, NICU: Neonatal Intensive Care Unit, VLBW: Very Low birth weight, PDA: Patent Ductus Arteriosus, RDS: Respiratory Distress Syndrome.

Introduction

Globally, about 2.4–2.8 million neonates died in 2017 and during this period south Asia and Africa had the highest neonatal mortality rate [1,2]. The neonatal mortality rate in Afghanistan is one of the highest in the world and it reported by UNICEF in 2019, 37 deaths per 1000 live births [3,4]. Prematurity, defined as neonate delivered before 37 weeks of gestation, is one of the major causes (35%) of early neonatal mortality in Afghanistan [5,6]. Apart from high mortality, Prematurity is also an important cause of neuromotor disability. It has been estimated that 15 million preterm babies are born worldwide every year [2,6]. Birth weight less than 1500g is classified as low birth weight [6].

Apnea is defined as a cessation of breathing for 20 seconds or longer or a shorter pause accompanied by bradycardia (<100 beats per minute), cyanosis, or pallor. In practice, many apneic events in preterm infants are shorter than 20 seconds. Approximately, all infants born at ≤ 28 weeks gestation have apnea; beyond 28 weeks gestation, the proportion of apnea decreases, from 85% at 30 weeks gestation to 20% at 34 weeks gestation [7,8].

Apnea or cessation of respiration is an important cause of mortality and brain damage in immature babies especially those with a gestation of fewer than 34 weeks or weight less than 1500g [9]. It is due to the immaturity of the central nervous system (apnea of prematurity) or secondary to other causes such as metabolic disturbances [10,11]. The prevalence of apnea in neonates with very low birth weight who admitted to NICU of the Aditya Hospital of India was 44% [12]. In the NICU of Enugu State University Teaching Hospital, South-East Nigeria, the prevalence of apnea was reported 84%, 47%, and 15% in ELBW, VLBW, and LBW neonates respectively [13]. The prevalence of apnea in preterm infants increase with low gestational age and birth weight of the newborn infants [14,15].

Despite high neonatal mortality due to prematurity and its complications, there is not enough scientific information regarding the epidemiology of AOP in preterm newborns of Afghanistan.

The present study was carried out to find the prevalence rate and risk factors of AOP in preterm neonates who were admitted to the Neonatal Intensive Care Unit (NICU), French Medical Institute for Mothers and Children (FMIC) Hospital, Kabul City. Such

information is highly required for better neonatal care, generation of hypotheses, further researches, and health policymaking.

Methods

Study Design, Setting, and Population

We conducted an analytic cross-sectional study in the Neonatal Intensive Care Unit (NICU), French Medical Institute for Mothers and Children (FMIC) Hospital, Kabul City during July-December 2019. The study population comprised of in-born and out-born preterm neonates (infants up to 28 days old) with weights less than 2500g.

Inclusion and Exclusion Criteria

The neonates of with a gestational age of fewer than 37 weeks and birth weight less than 2500g were enrolled in this study. No cases were excluded from the study.

Sample Size and Sampling Strategy

We calculated the sample size by Stata 14. A hypothesized difference of two years with standard deviation of 3.5 between mean maternal age of study and control group was took into consideration. With an alpha error of 10% and power of 80%, the two-sample means test estimated a sample size of 74 neonates. During the study period, thirty-six premature neonates who had apnea of prematurity (Apnea group) and the other thirty-nine premature neonates without apnea of prematurity (Control group) were recruited by census sampling method.

Variables

Main variables are

- The birth weight of neonates in gram was determined by accurate balance during the first 24hr of life. According to the birth weight the neonates were classified as low birth weight (1500-2500g), very low birth weight (1000-1500g), and extremely low birth weight (less than 1000g).
- The gestational age of neonates in weeks was determined by Last Menstrual Period (LMP) or antenatal maternal ultrasound or New Ballard Score. Prematurity was defined as birth occurring at less than 37 weeks of gestational age and was classified as early (less than 32 weeks of gestation), moderate (32-34 weeks of gestation) and late (35-36 weeks of gestation).
- The Apgar score of 1st minute of birth was recorded. The neonates with Apgar score of less than 7 at 1st minute were accepted as perinatal asphyxia.
- The apnea was defined as a cessation of breathing for 20 seconds or longer or a shorter pause accompanied by bradycardia (<100 beats per minute), cyanosis, or pallor. Persistent of apnea attacks in preterm neonates with no other obvious causes or despite the correction of secondary causes was accepted as apnea of prematurity. Continuous cardio-respiratory monitor and clinical examination detected the attacks of apnea.
- Neonatal sepsis was diagnosed by the presence of three of the five criteria: elevated C-reactive protein, leukocytosis or leucopenia, elevated count of premature granulocytes, temperature instability and symptoms of infection.

- Neonatal anemia was defined as hemoglobin level of less than 13gr/dl.
- Respiratory distress syndrome (RDS) was diagnosed by the presence of at least two of the following clinical signs, as well as characteristic radiographic finding of the chest: tachypnea (>60/min), dyspnea with inspiratory subcostal or intercostal retractions, nasal flaring, expiratory grunting and cyanosis in room air. On chest radiography, homogenous lung disease, classically described as having a ground-glass reticulo-granular appearance with air bronchograms, was a pathognomonic sign of RDS.
- The diagnosis of NEC was based on a combination of clinical symptoms and signs, including feeding intolerance, abdominal distention and bloody stools, as well as radiological features, such as pneumatosis intestinalis, portal venous gas and pneumoperitoneum.
- PDA was diagnosed by echocardiography and intraventricular hemorrhage was detected by the cranial ultrasound.
- The maternal age in year, type of delivery (vaginal or cesarean), and parity was defined as the number of births with the gestational age of more than 20 weeks.
- In-born newborns were those delivered at the Obstetric Unit of FMIC hospital, whereas out-born babies were those delivered at other hospitals.

Ethical Consideration

This study was approved by the Department of Neonatology (Protocol no 5 dates 3/2/2021), Kabul University Medical Science, and the Ethical Board of FMIC hospital. Although, this was an observational study and the data were obtained from the patient documents, verbal consents were taken from patients' guardians for participating in this study. The Helsinki Declaration was taken into consideration and all the personal information remained anonymous.

Data Collection Tools & Statistical Analysis

Initially, raw data were collected in data collection sheets and then entered in SPSS 24 software for statistical analysis. We detected the significance level by using independent t-test, Chi-square test and Binary Logistic Regression. A power of 80% and an alpha error of 0.1 were accepted so the p-value less than 0.1 was significant.

Results

Seventy-five preterm newborn babies were evaluated for the prevalence and pattern of AOP at the Neonatal Intensive Care Unit (NICU) of French Medical Institute for Mothers and Children (FMIC) Hospital, Kabul City. The demographic characteristics of the mentioned preterm neonates are described in Table-1. As shown in Table-2 the overall prevalence rate of AOP in newborn babies was 48% and the highest prevalence rates were seen within groups of ELBW and early preterm infants.

Table 1: Demographic Characteristics of the Participants.

Variables	Mean (SD)	Number (%)
Age in days	7.9 (2.6)	-
Birth weight in grams	1285.4 (259.6)	-
Gestational age in weeks	31.7 (2.4)	-
Apgar Scores of 1 st minute	7 (3)	-
Maternal Age in years	25.7 (4.2)	-
Number of Deliveries	2.4 (1.4)	-
Class of Birth Weight		
LBW (1500-2000 g)	-	11 (14.7)
VLBW (1000-1499)	-	57 (76)
ELBW (less than 1000g)	-	7 (9.3)
Class of GA		
Late Preterm (35-36 W)	-	14 (18.7)
Moderate Preterm (32-34 W)	-	25 (33.3)
Early Preterm (<32W)	-	36 (48)
Sex		
Boys	-	35 (46.7)
Girls	-	40 (53.3)
Mode of Delivery		
Vaginal	-	37 (49.3)
Cesarean	-	38 (50.7)
Place of Delivery		
In-born	-	62(82.7)
Out-born	-	13(17.3)

Table 2: Prevalence Rate of AOP in Different Categories of Preterm Neonates.

Categories of Neonates	Neonates with AOP, n	Neonates Without AOP, n	Prevalence Rate, %
All Neonates, n=75	36	39	48
Classes of Birth Weight, No			
LBW (1500-2000 g)	4	7	36.4
VLBW (1000-1499)	27	30	47.4
ELBW (less than 1000g)	5	2	71.4
Classes of Gestational Age, No			
Late Preterm (35-36 W)	5	9	35.7
Moderate Preterm (32-34W)	11	14	44
Early Preterm (<32W)	20	16	55.6
Sex, No			
Boys	19	16	54.3
Girls	17	23	42.5

After the comparison of data between apnea and non-apnea groups, as shown in (Table 3), the lower mean neonatal birth weight and anemia, as well as mean maternal age had a significant association with the development of AOP. The neonatal age, sex, gestational age, number of deliveries, and mode of delivery, as well as Sepsis, Perinatal Asphyxia, RDS, NEC, PDA, IVH, Hypocalcemia, Hyponatremia, Hypomagnesemia and hypokalemia had no statistically significant relationship with the risk of apnea in such infants. However, the risk of mortality was higher in the non-apnea group but such finding was not statistically significant.

Discussion

The current analytic cross-sectional study revealed the prevalence and figures of AOP in preterm neonates admitted to the NICU of the French Medical Institute for Mothers and Children Hospital, Kabul City. Based on the results, the overall prevalence rate of AOP was 48% and the highest rates were demonstrated within groups of ELBW and early preterm neonates, 71.4% and 55.6%

Table 3: Comparison of Factors Associated with AOP in Preterm Neonates.

Variables	Neonates with AOP (n=36)	Neonates Without AOP (n=39)	P-value (90%CI)
Birth Weight in grams, Mean (SD)	1233.33 (235.25)	1333.46 (274.44)	0.09 (-198 _ -1.4)
Gestational age in weeks, Mean (SD)	31.81 (2.01)	31.62 (2.73)	0.7 (-0.74_ 0.15)
Maternal Age in years, Mean (SD)	24.78 (3.68)	26.62 (4.58)	0.06 (-3.44 _ -0.23)
Number of Delivery, Mean (SD)	2.44 (1.3)	2.69 (1.5)	0.4 (-0.78_ 0.28)
Class of Birth Weight, No (%)			
LBW (1500-2000 g)	4 (36.4)	7 (63.6)	0.34
VLBW (1000-1499)	27 (47.4)	30 (52.6)	0.14
ELBW (less than 1000g)	5 (71.4)	2(28.6)	0.22
Class of GA, No (%)			
Late Preterm (35-36 W)	5 (35.7)	9 (64.3)	0.56
Moderate Preterm (32-34 W)	11 (44)	14 (56)	0.83
Early Preterm (<32W)	20 (55.6)	16 (44.4)	0.29
Sex, No (%)			
Boys	19 (54.3)	16 (45.7)	
Girls	17 (56.7)	23 (43.3)	
Odds Ratio (OR)	1.6	0.6	0.3
Mode of Delivery, No (%)			
Vaginal	18 (48.6)	19 (51.4)	
Cesarean	18 (47.4)	20 (52.6)	
Odds Ratio (OR)	1.05	0.95	0.45
Associate Illnesses No (%)			
Sepsis	12 (33.3)	12 (30.8)	0.5
Perinatal Asphyxia	10 (27.8)	11(28.2)	0.64
RDS	16 (44.4)	18 (46.1)	0.53
Anemia	8 (22.2)	4 (10.2)	0.05
NEC	4 (11.1)	5 (12.8)	0.55
PDA	4 (11.1)	9 (25)	1.14
IVH	3(8.3)	3 (7.8)	0.62
Hypocalcemia	9 (25)	14 (35.9)	0.22
Hyponatremia	4 (11.1)	6 (15.4)	0.42
Hypomagnesemia	4 (11.1)	5 (12.8)	0.55
Hypokalemia	11 (50.6)	7 (17.9)	0.12
Mortality, No (%)	12 (33.3)	9 (23.1)	
Risk Ratio (RR)	1.4	0.6	0.3

SD: Standard Deviation, GA: Gestational Age, g: Gram, W: Week, CI: Confidence Interval, LBW: Low Birth Weight, VLBW: Very Low Birth Weight, ELBW: Extremely Low Birth Weight, RDS: Respiratory Distress Syndrome, NEC: Necrotizing Enterocolitis, PDA: Patent Ductus Arteriosus, IVH: Intraventricular Hemorrhage.

respectively. The apnea of prematurity was more prevalent in boys than girls (Table 2). The current study also found a relationship of low neonatal birth weight and anemia as well as low maternal age with the development of AOP in neonates (Table 3). It means a lower neonatal birth weight and hemoglobin, as well as a younger maternal age increase the risk of AOP in newborn babies.

Shinde R et al. conducted a retrospective observational study on VLBW neonates admitted to NICU of Aditya Hospital, Hyderabad of India. According to the result of this study, the prevalence of AOP in very low birth weight was 44% [12]. Chidiebere et al. were carried out a prospective study at the Neonatal Intensive Care Unit (NICU) of Enugu State University Teaching Hospital, South-East Nigeria. The prevalence of AOP was demonstrated 20% and 14% in VLBW and LBW neonates respectively [13]. Henderson-Smart reported an incidence of 85% in premature infants born at 30 weeks, with a subsequent decrease to 20% in those born at 34 weeks [15]. Compared to the studies of Shinde R et al., Chidiebere et al. and Henderson-Smart et al., the current study found higher prevalence rates of AOP within groups of LBW and VLBW

premature infants (Table 2). This difference may be attributed to the higher prevalence rate of prematurity and associated factors of apnea in preterm neonates of our country.

Ogunlesi et al. conducted a retrospective study at the Olabisi Onabanjo University Teaching Hospital of Nigeria. This research found a significant relationship between neonatal birth weight and prevalence of AOP in newborn infants [16]. The association of fewer birth weights with AOP was also detected in our study. Therefore, the findings of both studies are similar. The current study also uncovered an association of young maternal age with AOP (Table 3). Fuchs et al. performed a large cohort study and revealed that younger women (20–24 years) had an increased risk of premature deliveries (aOR=1.08, 95% CI 1.01–1.15) [17]. As the younger maternal age is a risk factor of preterm birth, it may also explain the increased risk of apnea in such an infant group.

The association of neonatal anemia with AOP was the other important finding of the current study. Anemia causes a decrease in oxygen carrying capacity, which can lead to decreased oxygen

delivery to the central nervous system and a decrease in the efferent output of respiratory neuronal network, all of which raise the risk of apnea [18].

Conclusion

The current study demonstrated an overall prevalence rate of 48% for AOP in neonates and the highest rates appeared within groups of extremely low birth weight and early preterm newborns babies. The lower mean neonatal birth weight and neonatal anemia, as well as the younger maternal age were the risk factors of AOP in neonates. However, the incidence of mortality was higher in the non-apnea group; such finding was not statistically significant. These findings are most useful for better neonatal care, the generation of hypotheses, further researches, and health policymaking.

Acknowledgement

We gratefully acknowledge the French Medical Institute for Mother and Children Hospital for the cooperation in this study.

References

1. Hug L, Alexander M, You D, et al. National, regional, and global levels and trends in neonatal mortality between 1990 and 2017, with scenario-based projections to 2030: A systematic analysis. *The Lancet Global Health*. 2019; 7: 710-720.
2. Hug L, Sharrow D, You D. Levels and Trends in Child Mortality. UNICEF. 2017.
3. Neonatal mortality. UNICEF DATA. 2018.
4. Health. UNICEF Afghanistan. 2019.
5. Ansari N. The causes of neonatal mortality in Afghanistan. 48 th International course in health development. Amsterdam. 2012.
6. Kliegman RM, ST Geme JW, Blum NJ, et al. The High Risk Infants. *Nelson Textbook of Pediatrics*. U.S.A. Lange. 2020; 3865-3907.
7. Eichenwald EC. Apnea of Prematurity. *PEDIATRICS the Official Journal American Academy of Pediatrics*. 2016; 137: 20153757.
8. Dobson NR, Thompson MW, Hunt CE. Apnea. *Averys Neonatology*. Wolters Kluwer. 2018; 389-391.
9. Zhao J, Gonzalez F, Mu D. Apnea of prematurity: from cause to treatment. *Eur J Pediatr*. 2011; 170: 1097-1105.
10. Mishra S, Agarwal R, Jeevasankar M, et al. Apnea in the newborn. *Indian J Pediatr*. 2008; 75: 57-61.
11. Zhao J, Gonzalez F, Mu D. Apnea of prematurity: from cause to treatment. *European Journal of Pediatrics*. 2011; 170: 1097-1105.
12. Shinde R, Haridas K, Nagar P, et al. A study of survival of very low birth weight neonates in a tertiary care hospital. *International Journal of Contemporary Pediatrics*. 2019; 6: 857-862.
13. Chidiebere ODI, Uchenna E, Ikenna KN, et al. The Low-birth weight Infants: Pattern of Morbidity and Mortality in a Tertiary Healthcare Facility in the South Eastern Nigeria. *Annual of Medical and Health Science Research*. 2018; 8: 4-10.
14. Eric C. Eichenwald, Kristi L. Watterberg, Susan Aucott, et al. Apnea of Prematurity. *Pediatrics*. 2016; 137: 20153757.
15. Henderson-Smart DJ. The effect of gestational age on the incidence and duration of recurrent apnea in newborn babies. *Journal of Pediatric and Child health*. 1981; 17: 273-276.
16. Ogunlesi TA, Ogunfowora OB. Pattern and determinants of newborn apnea in an under-resourced Nigerian setting. *Nigerian Journal of Clinical Practice*. 2012; 15: 159-164.
17. Fuchs F, Monet B, Ducruet T, et al. Effect of maternal age on the risk of preterm birth: A large cohort study. *PLoS One*. 2018; 13: 0191002.
18. Kattwinkel J. Neonatal apnea: pathogenesis and therapy. *J Pediatr*. 1977; 90: 342-347.