

# Cyclic Hydrostatic Compress Force Regulates Apoptosis of Meniscus Fibrochondrocytes via Integrin $\alpha 5\beta 1$

Yang ZHANG<sup>1\*</sup>, Fazheng WANG<sup>2\*</sup>, Liangxiao BAO<sup>1</sup>, Jing LI<sup>1</sup>, Zhanjun SHI<sup>1</sup>, Jian WANG<sup>1</sup>

\*These authors contributed equally to this work.

<sup>1</sup>Department of Orthopedic Surgery, Nanfang Hospital, Southern Medical University, Guangzhou, China, <sup>2</sup>Department of Orthopedic Surgery, The First Hospital of Ka Shi, Kashi Xinjiang, China

Received November 21, 2018

Accepted April 4, 2019

Epub Ahead of Print June 6, 2019

## Summary

Meniscus is a semilunar fibrocartilaginous tissue, serving important roles in load buffering, stability, lubrication, proprioception, and nutrition of the knee joint. The degeneration and damage of meniscus has been proved to be a risk factor of knee osteoarthritis. Mechanical stimulus is a critical factor of the development, maintenance and repair of the meniscus fibrochondrocytes. However, the mechanism of the mechano-transduction process remains elusive. Here we reported that cyclic hydrostatic compress force (CHCF) treatment promotes proliferation and inhibits apoptosis of the isolated primary meniscus fibrochondrocytes (PMFs), via upregulating the expression level of integrin  $\alpha 5\beta 1$ . Consequently, increased phosphorylated-ERK1/2 and phosphorylated-PI3K, and decreased caspase-3 were detected. These effects of CHCF treatment can be abolished by integrin  $\alpha 5\beta 1$  inhibitor or specific siRNA transfection. These data indicate that CHCF regulates apoptosis of PMFs via integrin  $\alpha 5\beta 1$ -FAK-PI3K/ERK pathway, which may be an important candidate approach during meniscus degeneration.

## Key words

Integrin  $\alpha 5\beta 1$  • Meniscus • Cyclic hydrostatic compress force

## Corresponding author

J. Wang, Department of orthopedic surgery, Nanfang Hospital, Southern Medical University, 1838 Guangzhou Avenue, Guangzhou, Guangdong, 510515, China. E-mail: gjzy125@126.com

## Introduction

Menisci are semilunar fibrocartilaginous tissues, which buffer load on the knee joint, including load

transmission and shock absorption (Walker and Erkman 1975). They also serve important roles in knee joint stability, lubrication, proprioception, and nutrition of the articular cartilage (Fox *et al.* 2015). In 2015, it was estimated that there are approximately 1.7 million surgeries performed for menisci injuries worldwide and this number is rising rapidly (Merriam and Dunn 2015). There are different injury patterns between different populations (Tandogan *et al.* 2004): acute tears due to trauma are predominantly found in young people, while degenerative tears are found mainly in older people (Englund *et al.* 2009) associated with aging (Aufderheide and Athanasiou 2004). The majority of aging-related meniscus tears are unsuitable for repair treatments (Rai *et al.* 2013), which often require partial or complete removal of the meniscus, namely meniscectomy. However, although meniscectomy helps to relieve pain and improve function, it does not protect against the development of osteoarthritis (Hall *et al.* 2014). It has been proved that by removing the meniscus, the average stress of the knee can be increased by 3 folds, with an even greater magnitude of the peak stress (Krause *et al.* 1976). Particularly, the aging-related meniscal injury has been reported to be risk factors of knee osteoarthritis (Englund *et al.* 2012). Thus it is critical to understand the initial stage of the meniscal degeneration.

The meniscus cells are described as fibrochondrocytes (Makris *et al.* 2011). In nature, they experience a combination of dynamic and static stress, including compressive, shear and tensile forces (Abdelgaiad *et al.* 2015). Sufficient mechanical stimuli

play a critical role in maintaining the development, growth and functions of meniscus cells (McNulty and Guilak 2015). However, the transduction of mechanical signals into biology changes in these cells remains to be elusive.

Recent researches have begun to elucidate the role of integrin during this process. Integrin is a family of cell-surface molecules, responsible for extracellular-intracellular signaling transduction. Each integrin molecule is a heterodimer of  $\alpha$  and  $\beta$  subunits. There have been 18  $\alpha$  and 8  $\beta$  subunits identified in mammals (Humphries 2000), which are expressed in a tissue-specific manner in humans and mammals. Multiple integrins had been proved to be playing a role during mechano-transduction and sensing microenvironment in cartilage, however these integrins seem to act oppositely or complementarily. Among them, integrin  $\alpha 5\beta 1$  is a classic receptor for fibronectin (Woods Jr *et al.* 1994, Wright *et al.* 1997) in some tissues that are constantly exposed to mechanical stimuli, such as bladder smooth muscle, and nucleus pulposus of intervertebral disc (Xia and Zhu 2011).

Being consisted of fibrochondrocytes, menisci are structurally distinct from either the limb growth plate or the articular cartilage. They are unique in patterns of the cellular organization and antigenicity (Fox *et al.* 2015). The function of integrin  $\alpha 5\beta 1$  has not been identified during mechano-transduction in the meniscus, which is also a load transmission tissue. Here, in this study, we reported that cyclic hydrostatic compress force (CHCF) could promote cell proliferation and inhibit cell apoptosis in isolated primary meniscus fibrochondrocytes (PMFs) in vitro, via upregulating integrin  $\alpha 5\beta 1$  and the phosphorylation level of its downstream molecules, FAK, PI3K and ERK.

## Methods

### *Isolation and cell culture of primary meniscus fibrochondrocytes (PMFs)*

The use of animals for this study was approved by the Animal Care Council of Nanfang Hospital. Menisci of 12-week Sprague-Dawley rats were harvested and cut into small pieces. Fibrochondrocytes were released by digestion with 0.22 % (w/v) Type II collagenase (Sigma-Aldrich) and 0.25 % trypsinase (Sigma-Aldrich) in PBS containing 100 mg/ml streptomycin and 100 U/ml penicillin (Sigma-Aldrich) for 30 min at 37 °C. The supernatant was removed and

the remaining tissue was digested with Type II collagenase and trypsinase solution for an additional 3 hours and passed through a nylon cell strainer (70 mm, Corning). After rinsed with PBS for three times and prepared as a single cell suspension, cells were resuspended in growth medium of DMEM supplemented with 10 % FBS and 1 % penicillin/ streptomycin.  $4 \times 10^6$  cells were then seeded in a 60mm-dish and incubated in a humidified atmosphere at 37 °C and 5 % CO<sub>2</sub>. For experiments, the PMFs were then passaged and seeded into 6-well plates in triplicates, at the density  $1 \times 10^6$  cells/well.

### *Identification of PMFs*

For immunochemistry staining, cells were fixed in 4 % formalin solution for 15 min and permeabilized with 0.3 % Triton X-100 in PBS for 10 min. The cells were then incubated with primary antibodies, including anti-Type-I collagen (ab34710, Abcam, 1:500) and anti-Type-II collagen (ab34712, Abcam, 1:300). After washing with PBS, cells were incubated with secondary antibody labelled with fluorescence (A11012, Gibco).

### *Application of CHCF on cell culture*

CHCF was applied to cells by a computer-controlled pressure chamber (OTS, Taizhou, China), which allows sterile manipulation and up to 150 kPa of hydrostatic compress force. CHCF was applied on

**Table 1.** Primers used for qRT-PCR analysis

Target gene	Primer sequence (5'-3')
<i>Integrin <math>\alpha 1</math></i>	GATATTGGCCCTAACAGAC GCGATCGATTATTCCTC
<i>Integrin <math>\alpha 3</math></i>	GAATCACACCGAGGTCCACT GCATCTCCCCAGCCCGTTG
<i>Integrin <math>\alpha 4</math></i>	AAAGGCAGTACAAATCTATCC GAGCCCACCTAACATCAGTAAT
<i>Integrin <math>\alpha 5</math></i>	AGCGACTGGAATCCTCAAGACC AGTTGTTGAGTCCGTACCT
<i>Integrin <math>\alpha V</math></i>	TGTCAGCCCAGTCGTGTCTT GCTCAGCTCCGTGTCAATTC
<i>Integrin <math>\beta 1</math></i>	GGAGAAAATGTGATGCCATACAT TGGGCTGGTACAGTTTGTCA
<i>Integrin <math>\beta 3</math></i>	CACAACACGCACCGACACCT CCCCGGTTGAACCTCTTACACT
<i>GAPDH</i>	TGATTCTACCCACGGCAAGTT TGATGGGTTCCCATGATGA

confluent cells for experiments at the level of 150 kPa for 12 hours, and then removed. It had been confirmed that the pH of the growth medium was constant at 7.5 and the temperature was maintained at 37 °C.

#### Cell proliferation assay

The proliferation of the PMFs was detected over a seven-day period using CCK-8 solution according to the manufacturer's instructions. All experiments were performed in triplicates at least three times and representative results are shown.

#### qRT-PCR

Total RNAs were isolated using Trizol according to the manufacture instructions, and then reversed transcribed using iScript cDNA Synthesis Kit and amplified by PCR (SYBR green) using primers for each integrin subunit (Ma *et al.* 2016, Wei *et al.* 2014) (Table 1).

qRT-PCR was assayed with Applied Biosystems® 7500 machine. Normalization of samples was achieved by measurement of the endogenous GAPDH. All reactions were run in triplicates. A melting curve analysis was performed after the final PCR cycle, in order to check the presence of non-specific PCR products or primer-dimers. Efficiency of amplification was determined by a relative standard curve derived from serial dilutions.  $2^{-\Delta\Delta CT}$  method was used to calculate the relative expression level.

#### Cell transfection

Specific siRNA for integrin  $\alpha 5$  and  $\beta 1$  were transfected by Lipofectamine™ RNAiMAX (Invitrogen) reagent, according to the manufacturer's protocol. For each transfection, qRT-PCR was used to evaluate the expression level of target gene. The control groups were transfected by scrambled siRNA.

#### Western blotting analysis


PMFs were collected and washed twice in PBS. They were then lysed in RIPA buffer with proteinase and phosphatases inhibitor cocktail (ThermoFisher), for 20 min at 4 °C and centrifuged at 15,000 x g for 30 min at 4 °C. The supernatant was collected and the protein concentration was determined by BCA Protein Assay Kit (Invitrogen). Equal amounts of protein (15 µg) were separated by 10 % SDS-PAGE and were transferred electrophoretically onto a PVDF membrane. Western blotting analysis was performed as standard protocol.

The primary antibodies used for western blotting

were anti-integrin  $\alpha 5$  (ab150361, Abcam), anti-integrin  $\beta 1$  (ab179471, Abcam), anti-FAK (sc-271195, Santa Cruz), anti-Pho-FAK (sc-81493, Santa Cruz), anti-PI3K (4249, CST), anti-Pho-PI3K (sc-1331, Santa Cruz), anti-ERK1/2 (9102, CST) and anti-Pho-ERK1/2 (4376, CST). The expression of  $\beta$ -actin (sc-1615, Santa Cruz) was used as an internal control.

#### Flow cytometry analysis

The percentage of apoptotic cells was evaluated by staining cells with Annexin V-FITC (BD Biosciences).  $1 \times 10^6$  PMFs were re-suspended in 1 x binding buffer, and then 5 µl of Annexin V-FITC was added and incubated for 15 min at room temperature in the dark. The samples were then examined using a BD Accuri™ C6 flow cytometer (BD Biosciences).



**Fig. 1.** Schematic graph of computer-controlled pressure cell culture system. **A)** Air compressor. **B)** Computer-controlled valve. **C)** Computer. **D)** Pressure sensor. **E)** Cell incubator. **F)** Pressure gauge.

#### Statistical analysis

All results were expressed as means  $\pm$  standard deviation (Mean  $\pm$  SD). Statistical analysis was performed using Students *t* test, and  $p < 0.05$  was considered as statistically significant.

## Results

#### Characterization of primary meniscus Fibrochondrocytes (PMFs)

In monolayer cell culturing, the PMFs exhibited the morphology of polygonal as fibroblasts (Fig. 1A). They were strongly positive in Type I collagen staining (Fig. 1B), and were weakly positive in Type II collagen staining (Fig. 1C).

### Cyclic Hydrostatic Compress Force (CHCF) Promotes Proliferation and Inhibits Apoptosis of PMFs

The cell viability was increased to around 1.5 folds with the presence of CHCF (Fig. 2A). The PMFs showed a high apoptotic rate at about 63.81% ± 4.93% under regular culturing, which was decreased to 48.92% ± 6.92% when treated with CHCF (Fig. 2B). Flowcytometric analysis demonstrated that CHCF treatment was able to decrease early apoptotic and late apoptotic cells (Fig. 2C-G). Such effects of CHCF can be abolished by cilengitide, the integrin inhibitor (Fig. 2A, B).

### CHCF increases integrin $\alpha 5$ and $\beta 1$ expression level of PMFs cultured in vitro

To identify which integrin subunits in cultured PMFs were changed by CHCF stimulation, the mRNA expression of integrin subunits  $\alpha 1$ ,  $\alpha 3$ ,  $\alpha 4$ ,  $\alpha 5$ ,  $\alpha v$ ,  $\beta 1$  and  $\beta 3$  were measured by qRT-PCR (Fig. 3A-G). It was shown that among the tested subunits, the mRNA of integrin  $\alpha 5$  and  $\beta 1$  were significantly increased by CHCF stimulation (Fig. 3D, F), whereas other integrin subunits were not significantly affected. Consistently, at protein level, Western blot analysis found that integrin  $\alpha 5$  and  $\beta 1$  expression were significantly increased by CHCF treatment (Fig. 3H).

### CHCF modulates downstream molecules of integrin $\alpha 5\beta 1$


To further understand the mechanism of integrin  $\alpha 5\beta 1$  pathway in PMFs, the cells were transfected with integrin  $\alpha 5$  and/or  $\beta 1$  siRNA. Compared with the controls, the expression of integrin  $\alpha 5$  and  $\beta 1$  were significantly suppressed, at both mRNA and protein levels (Fig. 4A-C). Consequently, the enhanced cell proliferation and inhibited apoptosis from CHCF treatment were not observed (Fig. 4D, E).

To investigate the downstream molecules that


take part in the mechano-transduction progress, Western blot analyses were performed to evaluate the expression levels of focal adhesion kinase (FAK) and phosphorylated-FAK (Pho-FAK). Similarly as the expression pattern of integrin  $\alpha 5$  and  $\beta 1$ , Pho-FAK was increased by CHCF (Fig. 4F). Consequently, Pho-PI3K and Pho-ERK1/2 were also increased. Instead, FAK, PI3K and ERK1/2 were not affected (Fig. 4F). The protein level of caspase-3 was also decreased by CHCF (Fig. 4F). These CHCF effects on protein expression and phosphorylation levels could be completely inhibited by the integrin  $\alpha 5\beta 1$  inhibitor or siRNA transfection (Fig. 4F).

## Discussion


The basic functions of the meniscus are to enable the complex movements of tibiofemoral articulation of the knee joint, protecting the articular cartilage. During these movements, mechanical forces, including compressive, shear and tensile stresses, are transmitted to the meniscus dynamically and cyclically. Mechanical stimuli have been established to be a major regulator of normal tissue morphology and function under physiological condition. Meanwhile, it is also an important determinant factor for cell fate during pathological processes (Kessler *et al.* 2001). The alteration of mechanical condition may lead to changes in osmotic pressure, streaming potentials and current, tissue pH, and hydrostatic pressure gradients, which can be sensed and responded by fibrochondrocytes of meniscus (Frank and Grodzinsky 1987, Mak 1986, Mow *et al.* 1984). Sufficient mechanical stimulus was important for preventing apoptosis (Pirttiniemi *et al.* 2004), and maintaining the metabolic activities including glycosaminoglycan (GAG) production and proteoglycan (PG) synthesis (Jung *et al.* 2014, Behrens *et al.* 1989,



**Fig. 2.** Characterization of primary meniscus fibrochondrocytes (PMFs). **A**) Morphology of PMFs in light micrograph. **B**) Immunochemistry staining of type I collagen. **C**) Immunochemistry staining of type II collagen. Scale bar=100  $\mu$ m.



**Fig. 3.** Cyclic Hydrostatic Compressive Force (CHCF) Promotes Proliferation and Inhibits Apoptosis of PMFs. **A**) CCK-8 proliferation assay showing CHCF treatment increases cell viability, which can be abolished by integrin inhibitor. **B**) CHCF treatment decreases cell apoptosis, which can be abolished by integrin inhibitor. **C-G**) Flowcytometric analysis to detect the apoptotic PMFs with or without CHCF treatment.



**Fig. 4.** CHCF Increases Integrin  $\alpha 5$  and  $\beta 1$  Expression Level of PMFs Cultured *in vitro*. **A-G**) qRT-PCR analysis of mRNA expression level of different integrin subunits under with or without CHCF treatment. **H**) Western blot analysis showing the protein level of integrin  $\alpha 5$  and  $\beta 1$  with or without CHCF treatment.


Jurvelin *et al.* 1989). However, due to different cell type and different tissue structure, the effects of CHCF on meniscus fibrochondrocytes have not been identified. In this study, to mimic the real mechanical stimuli, CHCF was applied at the level of 150kPa and the interval of 12 hours according to previous study (McNulty and Guilak, 2015).

Integrin is a family of transmembrane adhesion molecules, composed of both  $\alpha$  and  $\beta$  subunits. They have been proved to be the major cell-surface receptors for cell migration and adhesion (Widgerow 2013), involved in cell-extracellular matrix interaction. Growing evidence suggests that mechanical stimuli may be transduced by signaling pathways mediated by integrin, modulating various cellular functions, including cell survival, proliferation, gap junction and motility, and protein expression (Gerthoffer and Gunst 2001, Hood and Cheresh 2002). Exposure to mechanical stimuli has been found to be able to activate specific integrin family

members. The expression of integrin  $\alpha 1\beta 1$ ,  $\alpha 2\beta 1$ ,  $\alpha 3\beta 1$ ,  $\alpha 5\beta 1$ ,  $\alpha v\beta 3$  and  $\alpha v\beta 5$  have been identified in chondrocytes (Kurtis *et al.* 2003, Salter *et al.* 1992), which can be altered by mechanical stimulation, local microenvironment and pathological process such as osteoarthritis (Kim *et al.* 2003, Lucchinetti *et al.* 2004). Integrin  $\beta 1$  alone was reported to be involved in the upregulation of aggrecan mRNA and suppression of matrix metalloproteinase-3 (MMP-3) mRNA levels by dynamic stretching of monolayer chondrocytes (Millward-Sadler *et al.* 1999, Millward-Sadler *et al.* 2000). Meanwhile, blocking integrin  $\alpha 5\beta 1$  in articular chondrocytes abolishes chondrocyte responses to dynamic stretching (Holledge *et al.* 2008) and leads to enhanced cell apoptosis due to upregulated matrix metalloproteinase-13 (MMP-13). Indeed, the degeneration of meniscus has been reported to be an important risk factor of osteoarthritis. The cellular mechanisms accounting for these pathological changes

may be the alterations in metabolic activities, such as proliferative activity and proteoglycan synthesis (Pirttiniemi *et al.* 2004). Our Western blot and qRT-PCR analysis found that, the expression of integrin  $\alpha 5\beta 1$  decreased as the degeneration of the meniscus processed

(data not shown). This remains elusive as our study shows that CHCF is able to upregulated integrin  $\alpha 5\beta 1$  expression. One of the possibilities is that other factors, such as shear stress or aging related changes play a more critical role during degeneration.



**Fig. 5. CHCF Modulates Downstream Molecules of Integrin.**

**A**) mRNA expression level of Integrin  $\alpha 5$  after specific siRNA transfection. **B**) mRNA expression level of Integrin  $\beta 1$  after specific siRNA transfection. **C**) Protein expression level of Integrin  $\alpha 5$  and  $\beta 1$  after specific siRNA transfection. **D**) CCK-8 proliferation assay on transfected PMFs, showing CHCF treatment does not affect cell viability. **E**) CHCF treatment does not affect cell apoptosis on PMFs transfected with siRNA. **F**) Western blot analysis of downstream molecules of integrin, showing that CHCF alters the phosphorylation level of FAK, PI3K and ERK1/2.

In the present study, FAK-PI3K/ERK pathway was found to be involved in CHCF induced mechano-transduction. FAK is a cytoplasmic tyrosine kinase located in the focal adhesion complex, transducing signals from integrins (Lal *et al.* 2007, Wen *et al.* 2009). Consistently, in other cell types (Diercke *et al.* 2011, Hong *et al.* 2010), FAK acts as an upstream regulator of p-ERK1/2 upon mechanical stimulation, which will

translocate into the nucleus to active transcription (Ory and Morrison 2004). Specific inhibition of ERK1/2 activation could result in apoptosis of human chondrocytes (Shakibaei *et al.* 2001). Previous study suggested that the inhibited integrin-ligand interactions induced the conformational changes of the uncleaved caspase-3 molecule, followed by enhanced auto-cleavage and greater amount of active caspase-3 molecules

(Buckley *et al.* 1999). Once treated with CHCF, the apoptosis of PMFs was inhibited by activation of integrin (Fig. 2). On the other hand, FAK could also directly activate caspase by facilitating PI3K activation (Chen and Guan 1994, Kiyokawa *et al.* 1998, Sonoda *et al.* 2000). Consistent with these previous findings, the phosphorylation level of PI3K and ERK were elevated by CHCF treatment.

One of the limitations in this study is the monolayer culture system. It has been proved that a three dimensional (3D) culture system may perform better in simulating physiological microenvironments of chondrocytes (Grodzinsky *et al.* 2000, Sanz-Ramos *et al.* 2012). In the further study we will develop a 3D system to mimic the meniscus structure (Pingguan-Murphy *et al.* 2005). Furthermore, it is quite difficult to measure the exact mechanical loadings on meniscus during movements of the knee joint (Chen *et al.* 2018). We had tried to mimic the *in vivo* stress pattern. However, although it had been confirmed that the pH of the growth

medium was constant at 7.5, the major concern of the system applied in this study was the potential effect on cell culture medium. Thus the mechanical stimuli could be further modulated, including treatment duration, magnitudes of the stress and cyclic patterns.

In summary, the present study provides an insight into the role of integrin  $\alpha 5\beta 1$  during mechano-transduction of the PMFs. Further studies will be focused on different patterns of mechanical stress and culturing conditions.

## Conflict of Interest

There is no conflict of interest.

## Acknowledgements

This work was supported by the National Natural Science Foundation of China [81501904], Guangdong Provincial Medical Science Foundation [A2017192], and Natural Science Foundation of Xinjiang Province [2016D01C021].

## References

- ABDELGAIED A, STANLEY M, GALFE M, BERRY H, INGHAM E, FISHER J: Comparison of the biomechanical tensile and compressive properties of decellularised and natural porcine meniscus. *J Biomech* **48**: 1389-1396, 2015.
- AUFDERHEIDE AC, ATHANASIOU KA: Mechanical stimulation toward tissue engineering of the knee meniscus. *Ann Biomed Eng* **32**: 1163-1176, 2004.
- BEHRENS F, KRAFT EL, OEGEMA JR TR: Biochemical changes in articular cartilage after joint immobilization by casting or external fixation. *J Orthop Res* **7**: 335-343, 1989.
- BUCKLEY CD, PILLING D, HENRIQUEZ NV, PARSONAGE G, THRELFALL K, SCHEEL-TOELLNER D, SIMMONS DL, AKBAR AN, LORD JM, SALMON M: RGD peptides induce apoptosis by direct caspase-3 activation. *Nature* **397**: 534, 1999.
- BUSCHMANN MD, GLUZBAND YA, GRODZINSKY AJ, HUNZIKER EB: Mechanical compression modulates matrix biosynthesis in chondrocyte/agarose culture. *J Cell Sci* **108**: 1497-1508, 1995.
- CATERSON B, LOWTHER DA: Changes in the metabolism of the proteoglycans from sheep articular cartilage in response to mechanical stress. *Biochim Biophys Acta Gen Subj* **540**: 412-422, 1978.
- CHAI D, ARNER E, GRIGGS, D. GRODZINSKY A:  $\alpha v$  and  $\beta 1$  integrins regulate dynamic compression-induced proteoglycan synthesis in 3D gel culture by distinct complementary pathways. *Osteoarthr Cartil* **18**: 249-256, 2010.
- CHEN H-C, GUAN J-L: Association of focal adhesion kinase with its potential substrate phosphatidylinositol 3-kinase. *Proc Nat Acad Sci* **91**: 10148-10152, 1994.
- CHEN H-Y, PAN L, YANG H-L, XIA P, YU W-C, TANG W-Q, ZHANG Y-X, CHEN S-F, XUE Y-Z WANG L-X: Integrin alpha5beta1 suppresses rBMSCs anoikis and promotes nitric oxide production. *Biomed & Pharmacother* **99**: 1-8, 2018.
- DIERCKE K, KOHL A, LUX CJ, ERBER R: Strain-dependent up-regulation of ephrin-B2 protein in periodontal ligament fibroblasts contributes to osteogenesis during tooth movement. *J Biol Chem* **286**: 37651-37664, 2011.
- ENGLUND M, GUERMAZI, A. LOHMANDER SL: The role of the meniscus in knee osteoarthritis: a cause or consequence? *Radiol Clin* **47**: 703-712, 2009.

- ENGLUND M, ROEMER FW, HAYASHI D, CREMA MD, GUERMAZI A: Meniscus pathology, osteoarthritis and the treatment controversy. *Nat Rev Rheumatol* **8**: 412-419, 2012.
- FOX AJ, WANIVENHAUS F, BURGE AJ, WARREN RF, RODEO SA: The human meniscus: a review of anatomy, function, injury, and advances in treatment. *Clin Anat* **28**: 269-287, 2015.
- FRANK EH, GRODZINSKY AJ: Cartilage electromechanics – II. A continuum model of cartilage electrokinetics and correlation with experiments. *J Biomech* **20**: 629-639, 1987.
- GERTHOFFER WT, GUNST SJ: Invited review: focal adhesion and small heat shock proteins in the regulation of actin remodeling and contractility in smooth muscle. *J Appl Physiol* **91**: 963-972, 2001.
- GRAY ML, PIZZANELLI AM, GRODZINSKY AJ, LEE RC: Mechanical and physicochemical determinants of the chondrocyte biosynthetic response. *J Orthopaedic Res* **6**: 777-792, 1988.
- GRODZINSKY, AJ, LEVENSTON ME, JIN M, FRAN EH: Cartilage tissue remodeling in response to mechanical forces. *Ann Rev Biomed Engin* **2**: 691-713, 2000.
- HALL M, WRIGLEY TV, METCALF BR, CICUTTINI FM, WANG Y, HINMAN RS, DEMPSEY AR, MILLS PM, LLOYD DG, BENNELL KL: Do moments and strength predict cartilage changes following partial meniscectomy? *Med Sci Sports Exerc* **47**: 1549-1556, 2014.
- HOLLEDGE MM, MILLWARD-SADLER S, NUKI G, SALTER D: Mechanical regulation of proteoglycan synthesis in normal and osteoarthritic human articular chondrocytes—roles for  $\alpha 5$  and  $\alpha V\beta 5$  integrins. *Biorheology* **45**: 275-288, 2008.
- HONG S-Y, JEON Y-M, LEE H-J, KIM J-G, BAEK J-A, LEE J-C: Activation of RhoA and FAK induces ERK-mediated osteopontin expression in mechanical force-subjected periodontal ligament fibroblasts. *Mol Cell Biochem* **335**: 263-272, 2010.
- HOOD JD, CHERESH DA: Role of integrins in cell invasion and migration. *Nat Rev Cancer* **2**: 91-100, 2002.
- HUMPHRIES MJ: Integrin structure. *Biochem Soc Trans* **28**: 311-339, 2000.
- JUNG J-K, SOHN W-J, LEE Y, BAE YC, CHOI J-K, KIM J-Y: Morphological and cellular examinations of experimentally induced malocclusion in mice mandibular condyle. *Cell Tissue Res* **355**: 355-363, 2014.
- JURVELIN JI, KIVIRANTA I, SÄÄMÄNEN AM, TAMMI M, HELMINEN HJ: Partial restoration of immobilization-induced softening of canine articular cartilage after remobilization of the knee (stifle) joint. *J Orthop Res* **7**: 352-358, 1989.
- KELLY TAN, WANG CCB, MAUCK RL, ATESHIAN GA, HUNG CT: Role of cell-associated matrix in the development of free-swelling and dynamically loaded chondrocyte-seeded agarose gels. *Biorheology* **41**: 223-237, 2004.
- KESSLER D, DETHLEFSEN S, HAASE I, PLOMANN M, HIRCHE F, KRIEG T, ECKES B: Fibroblasts in mechanically stressed collagen lattices assume a “synthetic” phenotype. *J Biol Chem* **276**: 36575-36585, 2001.
- KIM SJ, KIM EJ, KIM YH, HAHN SB, LEE JW: The modulation of integrin expression by the extracellular matrix in articular chondrocytes. *Yonsei Med J* **44**: 493-501, 2003.
- KIYOKAWA E, HASHIMOTO Y, KURATA T, SUGIMURA H, MATSUDA M: Evidence that DOCK180 up-regulates signals from the CrkII-p130Cas complex. *J Biol Chem* **273**: 24479-24484, 1998.
- KRAUSE WR, POPE M, JOHNSON R, WILDER D: Mechanical changes in the knee after meniscectomy. *J Bone Joint Surg Am* **58**: 599-604, 1976.
- KURTIS MS, SCHMIDT TA, BUGBEE WD, LOESER RF, SAH RL: Inte-mediated adhesion of human articular chondrocytes to cartilage. *Arthritis Rheum* **48**: 110-118, 2003.
- LAL H, VERMA S, SMITH M, GULERIA R, LU G, FOSTER D, DOSTAL D: Stretch-induced MAP kinase activation in cardiac myocytes: differential regulation through  $\beta 1$ -integrin and focal adhesion kinase. *J Mol Cell Cardiol* **43**: 137-147, 2007.
- LEE DA, BADER DL: Compressive strains at physiological frequencies influence the metabolism of chondrocytes seeded in agarose. *J Orthop Res* **15**: 181-188, 1997.
- LUCCHINETTI E, BHARGAVA MM, TORZILLI PA: The effect of mechanical load on integrin subunits  $\alpha 5$  and  $\beta 1$  in chondrocytes from mature and immature cartilage explants. *Cell Tissue Res* **315**: 385-391, 2004.

- MA D, KOU X, JIN J, XU T, WU M, DENG L, FU L, LIU Y, WU G, LU H: Hydrostatic compress force enhances the viability and decreases the apoptosis of condylar chondrocytes through integrin-FAK-ERK/PI3K pathway. *Int J Mol Sci* **17**: 1847, 2016.
- MAK AF: Unconfined compression of hydrated viscoelastic tissues: a biphasic poroviscoelastic analysis. *Biorheology* **23**: 371-383, 1986.
- MAKRIS EA, HADIDI P, ATHANASIOU KA: The knee meniscus: structure-function, pathophysiology, current repair techniques, and prospects for regeneration. *Biomaterials* **32**: 7411-7431, 2011.
- MAUCK R, BYERS B, YUAN X, TUAN R: Regulation of cartilaginous ECM gene transcription by chondrocytes and MSCs in 3D culture in response to dynamic loading. *Biomech Model Mechanobiol* **6**: 113-125, 2007.
- MCNULTY AL, GUILAK F: Mechanobiology of the meniscus. *J Biomech* **48**: 1469-1478, 2015.
- MERRIAM AR, DUNN MG: Meniscus tissue engineering. In: *Regenerative Engineering of Musculoskeletal Tissues and Interfaces*. SP NUKAVARAPU, JW FREEMAN, CT LAURENCIN (eds), Woodhead Publishing, 2015, pp 219-237.
- MILLWARD-SADLER S, WRIGHT M, LEE H-S, NISHIDA K, CALDWELL H, NUKI G, SALTER D: Integrin-regulated secretion of interleukin 4: a novel pathway of mechanotransduction in human articular chondrocytes. *J Cell Biol* **145**: 183-189, 1999.
- MILLWARD-SADLER S, WRIGHT M, DAVIES L, NUKI G, SALTER D: Mechanotransduction via integrins and interleukin-4 results in altered aggrecan and matrix metalloproteinase 3 gene expression in normal, but not osteoarthritic, human articular chondrocytes. *Arthritis Rheum* **43**: 2091-2099, 2000.
- MOW VC, HOLMES MH, LAI WM: Fluid transport and mechanical properties of articular cartilage: a review. *J Biomech* **17**: 377-394, 1984.
- ORY S, MORRISON DK: Signal transduction: implications for Ras-dependent ERK signaling. *Curr Biol* **14**: R277-R278, 2004.
- PINGGUAN-MURPHY B, LEE D, BADER D, KNIGHT M: Activation of chondrocytes calcium signalling by dynamic compression is independent of number of cycles. *Arch Biochem Biophys* **444**: 45-51, 2005.
- PIRTTINIEMI P, KANTOMA T, SORSA T: Effect of decreased loading on the metabolic activity of the mandibular condylar cartilage in the rat. *Eur J Orthod* **26**: 1-5, 2004.
- RAI MF, PATRA D, SANDELL LJ, BROPHY RH: Transcriptome analysis of injured human meniscus reveals a distinct phenotype of meniscus degeneration with aging. *Arthritis Rheum* **65**: 2090-2101, 2013.
- SAH RL, KIM YJ, DOONG JY, GRODZINSKY AJ, PLAAS AH, SANDY JD: Biosynthetic response of cartilage explants to dynamic compression. *J Orthop Res* **7**: 619-636, 1989.
- SALTER D, HUGHES D, SIMPSON R, GARDNERSANZ-RAMOS P, MORA G, RIPALDA P, VICENTE-PASCUAL M, IZAL-AZCARATE I: Identification of signalling pathways triggered by changes in the mechanical environment in rat chondrocytes. *Osteoarthritis Cartilage* **20**: 931-939, 2012.
- SHAKIBAEI M, SCHULZE-TANZIL G, DE SOUZA P, JOHN T, RAHMANZADEH M, RAHMANZADEH R, MERKER H-J: Inhibition of mitogen-activated protein kinase kinase induces apoptosis of human chondrocytes. *J Biol Chem* **276**: 13289-13294, 2001.
- SONODA Y, MATSUMOTO Y, FUNAKOSHI M, YAMAMOTO D, HANKS SK, KASAHARA T: Anti-apoptotic role of focal adhesion kinase (FAK) Induction of inhibitor-of-apoptosis proteins and apoptosis suppression by the overexpression of FAK in a human leukemic cell line, HL-60. *J Biol Chem* **275**: 16309-16315, 2000.
- TANDOĞAN RN, TAŞER O, KAYAALP A, TAŞKIRAN E, PINAR H, ALPARSLAN B, ALTURFAN A: Analysis of meniscal and chondral lesions accompanying anterior cruciate ligament tears: relationship with age, time from injury, and level of sport. *Knee Surg Sports Traumatol Arthrosc* **12**: 262-270, 2004.
- WALKER PS, ERKMAN MJ: The role of the menisci in force transmission across the knee. *Clinical orthopaedics and related research*: 184-192, 1975.
- WEI T, LUO D, CHEN L, WU T, WANG K: Cyclic hydrodynamic pressure induced proliferation of bladder smooth muscle cells via integrin alpha5 and FAK. *Physiol Res* **63**: 127-134, 2014.
- WEN H, BLUME PA, SUMPIO BE: Role of integrins and focal adhesion kinase in the orientation of dermal fibroblasts exposed to cyclic strain. *Int Wound J* **6**: 149-158, 2009.

- WIDGEROW AD: Chronic wounds – is cellular ‘reception’ at fault? Examining integrins and intracellular signaling. *Int Wound J* **10**: 185-192, 2013.
- WOODS VL JR, SCHRECK PJ, GESINK DS, PACHECO HO, AMIEL D, AKESON WH, LOTZ M: Integrin expression by human articular chondrocytes. *Arthritis Rheum* **37**: 537-544, 1994.
- WRIGHT MO, NISHIDA K, BAVINGTON C, GODOLPHIN JL, DUNNE E, WALMSLEY S, JOBANPUTRA P, NUKI G, SALTER DM: Hyperpolarisation of cultured human chondrocytes following cyclical pressure-induced strain: Evidence of a role for  $\alpha 5\beta 1$  integrin as a chondrocyte mechanoreceptor. *J Orthop Res* **15**: 742-747, 1997.
- XIA M, ZHU Y: Fibronectin fragment activation of ERK increasing integrin  $\alpha 5$  and  $\beta 1$  subunit expression to degenerate nucleus pulposus cells. *J Orthop Res* **29**: 556-561, 2011.