



ORIGINAL RESEARCH PAPER

Ophthalmology

ASSOCIATION OF INTRAOCULAR PRESSURE CHANGES IN DIFFERENT DEGREES OF MYOPIA

KEY WORDS: Intraocular Pressure Changes, Myopia, mild myopia, Moderate myopia, High myopia.

Dr Shubhnav Jain*

Department Of Ophthalmology, Rukmani Birla Hospital, Jaipur
*Corresponding Author

Dr Rohan Jain

Department of Ophthalmology ,SMS Hospital ,Jaipur

ABSTRACT

Introduction: Myopia, also known as “short-sightedness” or “near-sightedness,” is an increasingly widespread condition around the world. Myopia is commonly defined as a spherical equivalent (SE) ≤ 0.5 dioptres (D) and develops mainly during childhood and early adulthood when excessive elongation of the eye causes images of distant objects to fall in front of the retina with the eye at rest, resulting in a blurred distance vision. Myopia can be classified according to age of onset, clinical variety, etiology, and degree. According to the degree, myopia can be classified as mild/low myopia (-0.25D to -2.75D); moderate/medium myopia (-3.00D to -6.00D) and high myopia (Above -6.00D). **Aim and Objective:** To investigate the intraocular pressure changes in the different degrees of myopia. **Material and Methods:** This study was a clinical study carried out at the department of ophthalmology, rukmani birla hospital, Jaipur. An informed consent was gotten from all the subjects who were part of the study. Clinical tests performed on the subjects include case history, visual acuity, external eye examination, ophthalmoscopy, static retinoscopy and subjective refraction. The subjects with myopia were grouped into three according to the degree of myopia. Their intraocular pressure was measured using the indentation tonometry. Data was collected and uploaded into the Statistical Package for Social Sciences (SPSS) version 21 and the one-way ANOVA was used to test the null hypotheses at 0.05% level of significance and 95% confidence interval. **Result:** The study included a Sample size of 100 Patients. The table showed that 50 (50.00%) of the subjects had mild myopia; 42 (42.00%) had moderate myopia and 8 (8.00%) had high myopia. IOP of 10 - 15 were 20 (20.00%); 16 -20, 26 (26.00%); 21 - 25, 4 (4.00%); 26 - 30, 0 (0.00%). Minimum, maximum and mean IOP values for low myopes were 10.3, 30.6 and 17.13 mmHg respectively. For medium myopes, the minimum, maximum and mean IOP values were 11, 22 and 16.63 mmHg respectively. For high myopes, the minimum, maximum and mean IOP values were 18.8, 22 and 20.15 mmHg respectively. SPSS data analysis using the one-way ANOVA at 0.05 level of significance showed that there was a significant difference [$P(0.036) < 0.05$] in the IOP values between the different degrees of myopia. **Conclusion:** In conclusion, there was a significant difference in the IOP of different degrees of myopia. Eye care practitioners are advised to closely monitor the IOP of their myopic patients. For young children, as the myopia progresses, the IOP should also be monitored.

INTRODUCTION

Myopia, also known as “short-sightedness” or “near-sightedness,” is an increasingly widespread condition around the world, particularly in East Asia. It has been predicted that by 2050, there will be approximately five billion people, equivalent to 49.8% of the world population, suffering from myopia. Myopia is commonly defined as a spherical equivalent (SE) ≤ 0.5 dioptres (D) and develops mainly during childhood and early adulthood when excessive elongation of the eye causes images of distant objects to fall in front of the retina with the eye at rest, resulting in a blurred distance vision. High myopia (SE ≤ 6.0 D) and pathologic myopia (pathological retinal changes secondary to high myopia) are more harmful than low-grade myopia, which may cause uncorrectable visual impairment or blindness, including sight-threatening diseases such as glaucoma, retinal detachment, and macular holes.¹

Myopia may also occur in association with some systemic diseases such as Cornelia de Lange syndrome, Down's syndrome, Ehlers Danlos syndrome, Weil-Marchesani syndrome etc. Ocular diseases associated with myopia include retinopathy of prematurity, gyrate atrophy, homocystinuria, Wagner's syndrome etc. Symptoms include eye fatigue, photophobia, blurry vision when looking at distant objects, headaches caused by eyestrain etc. Myopia can be classified according to age of onset, clinical variety, etiology, and degree. According to the degree, myopia can be classified as low myopia (-0.25D to -2.75D); medium myopia (-3.00D to -6.00D) and high myopia (Above -6.00D).²

Intraocular pressure (IOP) is the pressure within the eyeball occurring as a result of continuous production and drainage of the aqueous humour. This aqueous humour is produced by the capillaries of the non-pigmented layer of the ciliary processes to supply oxygen to the avascular lens and cornea.

Aqueous humour is composed of ascorbic acid, amino acids, bicarbonate, sodium, chloride, glucose, phosphate, potassium, pyruvate and lactate. Aqueous production is mediated by three mechanisms; ultrafiltration, active transport or active secretion and diffusion.³ Aqueous humour is drained away by the trabecular outflow system and the uveoscleral outflow system. The trabecular outflow system consists of collector channels and the Schlemm's canal while the uveoscleral outflow system consists of the aqueous veins and the episcleral veins. Normal intraocular pressure is often taken to be between 11 mmHg and 21 mmHg with mean IOP as 16 ± 2.5 mmHg.² Measurement of intraocular pressure can be done by using a tonometer. Measurements with an ideal tonometer should be accurate, reproducible and repeatable. An ideal tonometer should be easy to use, portable, standardized.⁴ Types of tonometry include applanation tonometry, indentation tonometry, rebound tonometry, contour matching, intraocular implantation of pressure sensors and transpalpebral phosphene induction. This study was carried out to investigate the intraocular pressure changes in the different degrees of myopia.

MATERIALS AND METHODS

This study was a clinical study carried out at the department of ophthalmology, Rukmani Birla Hospital, Jaipur. An informed consent was gotten from all the subjects who were part of the study. Clinical tests performed on the subjects include case history, visual acuity, external eye examination, ophthalmoscopy, static retinoscopy and subjective refraction. The subjects with myopia were grouped into three according to the degree of myopia. Their intraocular pressure was measured using the indentation tonometry. Data was collected and uploaded into the Statistical Package for Social Sciences (SPSS) version 21 and the one-way ANOVA was used to test the null hypotheses at 0.05% level of significance and 95% confidence interval.

RESULT

A total of 100 subjects were used for this study.

Table 1: Distribution of subjects among the different degrees of myopia.

Degree of Myopia	No. of patients	Percentage
Mild	50	50
Moderate	42	42
High	8	8
Total	100	100

Table 1 showed the distribution of the subjects according to the degree of myopia. The table showed that 50 (50.00%) of the subjects had low myopia; 42 (42.00%) had medium myopia and 8 (8.00%) had high myopia.

Table 2: Intraocular pressure distribution of subjects with low myopia.

IOP (mmHg)	No. of patients	Percentage
10-15	20	20
16-20	26	26
21-25	4	4
26-30	0	0
Total	50	50

Table 2 showed the distribution of intraocular pressure values of subjects with low myopia. The table showed that subjects with IOP of 10 - 15 were 20 (20.00%); 16 -20, 26 (26.00%); 21 - 25, 4 (4.00%); 26 -30, 0 (0.00%).

Table 3: Intraocular pressure distribution of subjects with medium myopia.

IOP (mmHg)	No. of patients	Percentage
10-15	22	22
16-20	20	20
21-25	0	0
26-30	0	0
Total	42	42

Table 3 showed the distribution of intraocular pressure values of subjects with medium myopia. The table showed that subjects with IOP of 10 - 15 were 22 (22.00%); 16 -20, 20 (20.00%); none of the subjects had IOP above 20mmHg.

Table 4: Intraocular pressure distribution of subjects with high myopia.

IOP (mmHg)	No. of patients	Percentage
10-15	0	0
16-20	8	8
21-25	0	0
26-30	0	0
Total	8	8

Table 4 showed the distribution of intraocular pressure values of subjects with high myopia. The table showed that none of the subjects had IOP of 10 - 15 and above 20 mmHg; 8 (8.00%) had IOP between 16 and 20 mmHg.

Table 5: Comparison of intraocular pressure values in the degrees of myopia.

Degree of Myopia	No. of patients	Min	Max	Mean	SD
Mild	50	10.3	30.6	17.13	3.97
Moderate	42	11	22	16.63	3.09
High	8	18.8	22	20.15	1.29

Table 5 showed the comparison of IOP values in the different degrees of myopia. The table showed that the minimum, maximum and mean IOP values for low myopes were 10.3, 30.6 and 17.13 mmHg respectively. For medium myopes, the minimum, maximum and mean IOP values were 11, 22 and 16.63 mmHg respectively. For high myopes, the minimum, maximum and mean IOP values were 18.8, 22 and 20.15 mmHg respectively. SPSS data analysis using the one-way ANOVA at 0.05 level of significance showed that there was a significant difference [P(0.036) < 0.05] in the IOP values

between the different degrees of myopia.

DISCUSSION

Myopia occurs mostly as a result of either an axial elongation of the eyeball or curvature increase of the cornea or lens. It is listed as one of the important risk factors in the development of lacquer cracks, subretinal neovascularization, primary open angle glaucoma, retinal detachment and dense cataract. Intraocular pressure plays a critical role in the pathogenesis of glaucoma and myopia. Elevated intraocular pressure causes an increase in the axial length. Edgar and Rudnicka⁵ found that low myopia was associated with a doubling of the risk of glaucoma at any age and a three-fold increase with medium and high myopia compared to those with emmetropia. The mean IOP values for the different degrees of myopia in this study were similar to related studies that also found a statistically significant change in intraocular pressure with different degrees of myopia. Manny et al.⁶ worked on the associations among intraocular pressure, ethnicity and refractive error and found that intraocular pressure differed among refractive error. Higher intraocular pressure was found more in children with low and medium myopia than in those with high hyperopia.

Sowjanya et al.⁷ found a positive correlation between intraocular changes with refractive errors. Tham et al.⁸ found a significant interaction between IOP and myopia in primary open angle glaucoma in a study on the joint effects of intraocular pressure and myopia on the risk of open angle glaucoma in a multi-ethnic Asian population. There are other works which do not agree with the results of this study. Chinawa et al.⁹ reported a lack of correlation between myopia and intraocular pressure and intraocular pressure and axial length. They however, reported a linear correlation between myopia and axial length. They also found that increased intraocular pressure causes scleral stress and creep resulting in elongation of the axial length and scleral stretch. A study by Lee et al.¹⁰ on intraocular pressure associations with refractive error and axial length in children did not support an association between IOP and refractive error or axial length. This change in intraocular pressure with the different degrees of myopia could also be attributed to reduced choroidal circulation and consequent oxidative stress. Perfusion of the retina, choroid and optic nerve head is critical to maintaining normal vision.

Samra et al.¹¹ investigated the change in subfoveal choroidal blood flow in patients with glaucoma and assessed the effect of myopia as one of the vascular risk factors for glaucoma on this flow. They concluded that subfoveal choroidal laser Doppler flowmetry parameters were reduced in patients with myopia and primary open-angle glaucoma than in nonmyopic glaucomatous patients with age-matched comparison. They suggested impaired choroidal circulation caused by myopia as an additional risk factor involved in glaucomatous damaging process. Sugiyama et al.¹² evaluated the blood circulation in the retina, peripapillary choroid and optic disc in normal -tension glaucoma and correlated them with visual field measurement, retinal vessel width, optic disc pallor and blood pressure. They found a possible association of a decrease in retinal, choroidal and optic disc blood flow in the inferior and nasal quadrant as well as vessel width in the superior quadrant and showed an increasing blood flow for enlargement of the pallor. A decrease in blood pressure was found to be related to blood flow choroid and optic disc impairment. Changes in intraocular pressure with the different degrees of myopia have been attributed to reduced ocular perfusion.

Zhi et al.¹³ found a reduction of retinal blood flow and optic nerve head perfusion with increasing intraocular pressure in rats. A study by Wang et al.¹⁴ to evaluate the peripapillary and parafoveal perfusion of young healthy myopic subjects with spectral domain optical coherence tomography angiography

found significant differences in the retinal flow index and vessel density in the peripapillary area but not parafoveal area among low and medium myopes. The high myopes had the lowest peripapillary retinal flow index and vessel density.

CONCLUSION

In conclusion, there was a significant difference in the IOP of different degrees of myopia. Eye care practitioners are advised to closely monitor the IOP of their myopic patients. For young children, as the myopia progresses, the IOP should also be monitored.

REFERENCES

1. M. A. Bullimore and N. A. Brennan, "Myopia control: why each diopter matters," *Optometry and Vision Science*, vol. 96, no. 6, pp. 463-465, 2019.
2. Khurana AK. *Comprehensive Ophthalmology*. 5th ed, New Delhi: New Age International Publishers.2012;32-256.
3. Stamper RL, Liebermann MF, Drake MV, Mosby: *Becker-Shaffer's Diagnosis and Therapy of the Glaucomas*. 8th ed. Houston: Elsevier.2009;6-62
4. Yolcu U, Ilhan A, Tas A. *Glaucoma - Intraocular pressure and Aqueous Dynamics*. 1st ed. Chester: CC Publishers.2016;44-54.
5. Edgar FD, Rudnicka RA. *Glaucoma Identification and Co-management*. Edinburgh: Butterworth Heinemann Elsevier.2007;7.
6. Manny RE, Mitchell L, Cotter SA, JonesJordan LA, Kleinstejn RN, Multi DO. Intraocular pressure, Ethnicity and Refractive Error. *Optom Vis Sci*.2011; 88(12):1445-1453.
7. Sowjanya M, Sathyavati K, Biradar K, Girish B. A study on the association of intraocular pressure changes with refractive errors in Bidar population. *Int J Current Res Aca Rev*.2015;3(4):320-327.
8. Tham Y, Aunt T, Fan Q, Saw S, Siantar RG, Wong TY, Cheung C. Joint effects of intraocular pressure and myopia on risk of Primary Open-Angle Glaucoma. *The Singapore Epidemiol of Eye Diseases Study Sci Rep*.2016;382-387.
9. Chinawa EN, Adio AO, Chukwuka OI. Is there a causal relationship between myopia and intraocular pressure? *Br J Med Med Res*.2017;20(10):1-7.
10. Lee SJ, Choi RY, Lee EJ, Choi YH, Lee SH, Oum SB. Relationship between Intraocular Pressure and Systemic Health Parameters in the Korean population. *Kor J Ophthalmol*.2002;16:13-19.
11. Samra WA, Pournaras C, Riva C, Emarah M. Choroidal hemodynamics in myopic patients without primary open angle glaucoma. *Acta Ophthalmol*.2013;91:371-375.
12. Sugiyama T, Schwartz B, Takamoto T, Azumal I. Evaluation of the circulation in the retina, peripapillary choroid and optic disk in normal-tension glaucoma. *Ophthalmol Res*.2000;32(2-3): 79-86.
13. Zhi Z, Cepurna OW, Johnson CE, Morrison CJ, Wang KR. Impact of intraocular pressure on changes of blood flow in the retina, choroid and optic nerve head in rats investigated by optical microangiography. *Biomed Opt Express*.2012;3(9): 220-223.
14. Wang X, Kong X, Jiang C, Li M, Yu J, Sun X. Is peripapillary perfusion related to myopia in healthy eyes? *BMJ*.2016;6:1-7.