

## *Ralstonia Solanacearum* Species Complex and Bacterial Wilt Disease

Fanhong Meng\*

Boyce Thompson Institute for Plant Research, 533 Tower Rd, Ithaca, NY 14853, USA

### Abstract

The bacterium *Ralstonia solanacearum* is the causal agent for bacterial wilt on more than 200 plant species from 50 botanical families, including important crops such as potato, tomato, eggplant, pepper, tobacco and banana. *R. solanacearum* is considered a species complex, and is also a widely accepted model organism for the study of bacterial pathogenicity in plants. This review discusses the disease caused by *R. solanacearum*, the classification of the pathogen, the major virulence and pathogenicity factors and their complex regulation in *R. solanacearum*.

**Keywords:** *Ralstonia solanacearum*; Bacterial wilt; Species complex; Classification; Virulence; Pathogenicity factors; Regulation.

### Introduction

The bacterium *Ralstonia solanacearum* is a widely accepted model organism for the study of pathogenicity in plants [1]. This soil-borne bacterium, which belongs to the betaproteobacteria, is responsible for bacterial wilt on more than 200 plant species from 50 botanical families, including important crops such as potato, tomato, eggplant, pepper, tobacco and banana. In fact, bacterial wilt is considered the single most destructive bacterial plant disease because of its extreme aggressiveness, wide geographic distribution, and unusually broad host range [2].

*R. solanacearum* inhabits the vascular tissue of its hosts. The bacterium normally invades plant roots from the soil through wounds or natural openings where secondary roots emerge [3], colonizes the intercellular space of the root cortex and vascular parenchyma, and eventually enters the xylem vessels and spreads up into the stem and leaves, where the pathogen cell density commonly surpasses  $10^9$  CFU/g of host tissue [3,4]. After *R. solanacearum* has colonized the xylem, large numbers of bacterial cells are shed from roots, providing a pathway for bacteria to return to the soil and initiate new infections [1]. Affected plants suffer chlorosis, stunting, wilting, and usually die rapidly.

### *R. solanacearum* species complex and classification

*R. solanacearum* is considered a species complex—a heterogeneous group of related but genetically distinct strains [5,6]. DNA-DNA hybridization studies have revealed that the identity between *R. solanacearum* genomes is often less than the 70% threshold level commonly expected within a bacterial species [7,8]. Gillings and Fahy first used the term “species complex” to describe the high genetic variation between isolates, and Taghavi et al. [9] expanded the concept of the *R. solanacearum* species complex by including two closely related species from Indonesia, *Ralstonia syzygii* (a pathogen of clove trees) and the agent of blood disease of banana, known as the BDB. *R. solanacearum* species complex strains are highly competent for genetic exchange *in planta* [10] and show substantial pathogenic variability in host range and aggressiveness [5]. However, the genetic basis for this variation is unknown. For the past four decades, two different systems, race and biovar, have been widely used to differentiate *R. solanacearum* strains [11–13]. Historically, *R. solanacearum* was subdivided into “races” based loosely on host range [13]. Race 1 strains attack tobacco, many other solanaceous crops and many hosts in other plant families;

race 2 strains are limited to musaceous species including *Heliconia* spp. and triploid banana; race 3 strains primarily attack potato; race 4 strains are particularly virulent on ginger; race 5 strains infect and cause disease on mulberry tree, and are only found in China [14]. The race structure of *R. solanacearum* is poorly defined and not taxonomically useful.

Based on their different abilities to utilize and oxidize several disaccharides (cellobiose, lactose, and maltose) and hexose alcohols (dulcitol, mannitol, and sorbitol), *R. solanacearum* strains were originally divided into five biovars [12]. Biovar 1 strains metabolize none of them; biovar 2 strains only metabolize disaccharides; biovar 3 strains metabolize all of them; biovar 4 strains metabolize only hexose alcohols; biovar 5 strains metabolize all of them except dulcitol and sorbitol [12,14]. Later, a new group of *R. solanacearum* isolates from the Amazon basin was differentiated from original biovar 2 using ribose and trehalose [15]. This group is named biovar 2-T or biovar N2 and the original biovar 2 strains are now referred to as 2-A. Except for biovar 2-A, which almost always corresponds to race 3, and biovar 5, which is identical to race 5, there is no correlation between biovars and races.

Recently, a new phylogenetic classification system consisting of four phylotypes was formed based on phylogenetic analysis of sequence data from the 16S-26S internal transcribed spacer region, the *egl* gene, the *hrpB* gene, and the *mutS* gene [5,6,16]. The phylotypes correlate with the geographical origin of the strains: phylotype I includes strains originating primarily from Asia, phylotype II from America, phylotype III from Africa and surrounding islands in the Indian Ocean, and phylotype IV from Indonesia [5,6]. Each phylotype is further divided into sequevars, which are defined as isolates with less than 1% nucleotide variation in the endoglucanase (*egl*) locus [6,16]. The *R. solanacearum* phylotype system is a stable and phylogenetically meaningful classification scheme that represents the evolutionary lineages within the species complex [6].

**\*Corresponding author:** Fanhong Meng, Boyce Thompson Institute for Plant Research, 533 Tower Rd, Ithaca, NY 14853, USA, Tel: (607) 220 9610; Email: [fm352@cornell.edu](mailto:fm352@cornell.edu)

**Received** February 26, 2013; **Accepted** March 19, 2013; **Published** March 26, 2013

**Citation:** Meng F (2013) *Ralstonia Solanacearum* Species Complex and Bacterial Wilt Disease. J Bacteriol Parasitol 4: e119. doi:10.4172/2155-9597.1000e119

**Copyright:** © 2013 Meng F. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## The Major Virulence and Pathogenicity Factors in *R. solanacearum* and Their Regulation

### The major virulence and pathogenicity factors

Many factors contribute to virulence and pathogenicity of *R. solanacearum* strains, including a heterogenous polymer of N-acetylated extracellular polysaccharide I (EPS I) [17-19], the type III secretion system (T3SS) [20], flagella-driven swimming motility and type IV pili-driven twitching motility [21-24], Cell-Wall-Degrading Enzymes (CWDEs) and type II secretion system (T2SS) [25-27].

*R. solanacearum* EPS I-deficient mutants are nearly avirulent and had reduced stem colonization compared to wild type [3,28-30]. Two functions have been hypothesized for EPS I: one is physically blocking the vascular system and thereby altering water movement [19,28,29], the other is protecting *R. solanacearum* from plant antimicrobial defenses by cloaking bacterial surface features that could be recognized by hosts [3,30]. In *R. solanacearum*, the T3SS is encoded by a 23-kb *hrp* gene cluster on the megaplasmid [20] and is essential for pathogenicity, since *R. solanacearum* *hrp* mutants are unable to cause disease in many susceptible plants [31-33]. Both flagella-driven swimming motility and type IV pili-driven twitching motility contribute to the ecological fitness and virulence of *R. solanacearum* [21-24]. While swimming motility contributes to virulence only in the early stage of host colonization and invasion [21,22,34], the type IV pili and twitching motility are important for several stages of wilt disease development [23,24]. *R. solanacearum* secretes several plant CWDEs via the T2SS that contribute to the pathogen's ability to cause wilt, although none is essential for disease [25-27]. A T2SS mutant was much less virulent than the mutant lacking all known CWDEs, suggesting that additional extracellular proteins secreted by the T2SS contribute to the virulence of *R. solanacearum* [27].

### Complex regulation of virulence and pathogenicity factors

The key virulence and pathogenicity factors of *R. solanacearum* are controlled by a complex regulatory network that responds to multiple signals [35]. The central player of this regulatory network is the transcriptional regulator PhcA, involved in the Phc cell density sensing system [36]. PhcA positively regulates the production of EPS I, Pme and Egl exoproteins, and an acyl-homoserine lactone (AHL) quorum sensing system called SolI/R. PhcA also represses the expression of genes involved in motility, T3SS, polygalacturonase and siderophore production [24,37-42]. The levels of active PhcA protein are controlled by a diffusible endogenous signal molecule, 3-hydroxypalmitic acid methyl ester (3-OH PAME). When extracellular 3-OH PAME accumulates above threshold concentrations (i.e., at high cell density in a confined space, such as the plant vascular system), PhcA is expressed, resulting in activation or repression of its various target genes [35,43].

In addition to the Phc cell confinement sensing system mediated by 3-OH-PAME, *R. solanacearum* also produces acylated homoserine lactones (acyl-HSLs) and has a functional acyl-HSL-dependent autoinduction system mediated by the above-mentioned SolI-SolR regulators, which are LuxI/LuxR homologs [40]. At present, only one gene of unknown function (*aidA*) is known to be regulated by SolI-SolR in phylotype II seq 7 strain AW1. *aidA* is not present in strain GMI1000 [40]. While inactivation of *solIR* does not affect any virulence factor production, the role of SolIR in the physiology of *R. solanacearum* requires more investigation. The acyl-HSL-dependent autoinduction system in *R. solanacearum* is part of a more complex autoregulatory

hierarchy, since expression of *solR* and *soli* requires PhcA, which is itself controlled by another autoregulatory system that responds to 3-OH-PAME [40]. Expression of SolIR is also dependent on the alternate sigma factor RpoS [44].

*R. solanacearum* regulates deployment and use of its T3SS with a complex environmentally responsive signal transduction cascade via the Prh system [35,45]. *R. solanacearum* T3SS genes are induced by bacterial contact with plant cells or cell wall fragments [46]. This contact-dependent induction requires PrhA [47], an outer membrane-bound protein that may sense and transfer signal to activate T3SS expression through a linear pathway including the PrhR, PrhI, PrhJ, HrpG and HrpB proteins [48,49]. HrpG, an OmpR family two-component response regulator, is involved in *hrpB* gene activation in response to both nutrient and metabolic signals. HrpG plays a central role in regulating T3SS gene expression [49]. In addition, HrpG also independently regulates bacterial production of plant cell wall-degrading enzymes, exopolysaccharide, and the phytohormones ethylene and auxin [50]. Moreover, in culture the activity of HrpG is modulated by the Phc confinement sensing system [41]. HrpB is an AraC-like transcriptional activator required for the transcription of other *hrp* genes involved in T3SS and the effector protein genes [51-53]. HrpB was also found to regulate genes involving in chemotaxis, biosynthesis or catabolism of various low-molecular-weight chemical compounds, and siderophore production and uptake [51].

### Conclusion

The present review has focused on the bacterial wilt disease caused by *R. solanacearum*, the classification of the pathogen, the major virulence and pathogenicity factors in *R. solanacearum* and their complex regulation. *R. solanacearum*, as a model organism for the study of bacterial pathogenicity in plants, survives in diverse environments and cause destructive disease on a wide range of plants by precisely controlling its gene expression through an elaborate regulatory network. Over the years, significant progress has been made in studying *R. solanacearum* species complex and bacterial wilt disease. However, further research is needed to better understand questions such as how does *R. solanacearum* regulate its gene expression in different niches, especially inside host plants? How do different plants respond to *R. solanacearum*? A better understanding of these questions will help us to design innovative tools and strategies to combat bacterial wilt disease more effectively.

### References

1. Hayward AC (1991) Biology and epidemiology of bacterial wilt caused by *Pseudomonas solanacearum*. Annu Rev Phytopathol 29: 65-87.
2. Prior P, Allen C, Elphinstone JG (1998) Bacterial Wilt Disease: Molecular and Ecological Aspects. Springer.
3. Araud-Razou I, Vasse J, Montrozier H, Etchebar C, Trigalet A (1998) Detection and visualization of the major acidic exopolysaccharide of *Ralstonia solanacearum* and its role in tomato root infection and vascular colonization. European Journal of Plant Pathology 104: 795-809.
4. Vasse J, Frey P, Trigalet A (1995) Microscopic studies of intercellular infection and protoxylem invasion of tomato roots by *Pseudomonas solanacearum*. Mol Plant Microbe Interact 8: 241-251.
5. Allen C, Prior P, Hayward AC (2005) Bacterial Wilt Disease and the *Ralstonia solanacearum* species complex. APS.
6. Prior P, Fegan M (2005) Recent developments in the phylogeny and classification of *Ralstonia solanacearum*. Acta Hort 695: 127-136.
7. Palleroni NJ, Doudoroff M (1971) Phenotypic characterization and

- deoxyribonucleic acid homologies of *Pseudomonas solanacearum*. J Bacteriol 107: 690-696.
8. Hayward AC, Hartman GL (1994) Bacterial Wilt: The disease and its causative agent, *Pseudomonas solanacearum*. CAB International.
  9. Taghavi M, Hayward C, Sly LI, Fegan M (1996) Analysis of the phylogenetic relationships of strains of *Burkholderia solanacearum*, *Pseudomonas syzygii*, and the blood disease bacterium of banana based on 16S rRNA gene sequences. Int J SystBacteriol 46: 10-15.
  10. Bertolla F, Frostergard A, Brito B, Nesme X, Simonet P (1999) During infection of its host, the plant pathogen *Ralstonia solanacearum* naturally develops a state of competence and exchanges genetic material. Mol Plant Microbe Interact 12: 467-472.
  11. Buddenhagen I, Kelman A (1964) Biological and physiological aspects of bacterial wilt caused by *Pseudomonas solanacearum*. Annu Rev Phytopathol.
  12. Hayward AC (1964) Characteristic of *Pseudomonas solanacearum*. J ApplBact 27: 265-277.
  13. Buddenhagen I, Sequeira L, Kelman A (1962) Designation of races in *Pseudomonas solanacearum*. Phytopathology 52: 726.
  14. He YL, Sequeira L, Kelman A (1983) Characteristics of strains of *Pseudomonas solanacearum* from China. Plant Dis 67: 1357-1361.
  15. Hayward AC(1994) Systematics and phylogeny of *Pseudomonas solanacearum* and related bacteria. In: Hayward AC, Hartman GL (Eds.) Bacterial Wilt: The Disease and its Causative Agent, *Pseudomonas solanacearum*.Wallingford, CAB International.
  16. Poussier S, Prior P, Luisetti J, Hayward C, Fegan M (2000) Partial sequencing of the hrpB and endoglucanase genes confirms and expands the known diversity within the *Ralstonia solanacearum* species complex. Syst Appl Microbiol 23: 479-486.
  17. McGarvey JA, Denny TP, Schell MA (1999) Spatial-Temporal and Quantitative Analysis of Growth and EPS I Production by *Ralstoniasolanacearum* in Resistant and Susceptible Tomato Cultivars. Phytopathology 89: 1233-1239.
  18. Orgambide G, Montrozier H, Servin P, Roussel J, Trigalet-Demery D, et al. (1991) High heterogeneity of the exopolysaccharides of *Pseudomonas solanacearum* strain GMI 1000 and the complete structure of the major polysaccharide. J BiolChem 266: 8312-8321.
  19. Denny TP (1995) Involvement of bacterial polysaccharides in plant pathogenesis. Annu Rev Phytopathol 33: 173-197.
  20. VanGijsegem F, Gough C, Zischek C, Niqueux E, Arlat M, et al. (1995) The hrp locus of *Pseudomonas solanacearum*, which encodes the production of a type III secretion system, encodes eight proteins related to components of the bacterial flagellarbiogenesis complex. Mol Microbiol 15: 1095-1114.
  21. Tans-Kersten J, Brown D, Allen C (2004) Swimming motility, a virulence trait of *Ralstonia solanacearum*, is regulated by FlhDC and the plant host environment. Mol Plant Microbe Interact 17: 686-695.
  22. Tans-Kersten J, Huang H, Allen C (2001) *Ralstonia solanacearum* needs motility for invasive virulence on tomato. J Bacteriol 183: 3597-3605.
  23. Kang Y, Liu H, Genin S, Schell MA, Denny TP (2002) *Ralstonia solanacearum* requires type 4 pili to adhere to multiple surfaces and for natural transformation and virulence. Mol Microbiol 46: 427-437.
  24. Liu H, Kang Y, Genin S, Schell MA, Denny TP (2001) Twitching motility of *Ralstonia solanacearum* requires a type IV pilus system. Microbiology 147: 3215-3229.
  25. Huang Q, Allen C (2000) Polygalacturonases are required for rapid colonization and full virulence of *Ralstonia solanacearum* on tomato plants. PhysiolMol Plant Pathol 57: 77-83.
  26. Denny TP, Carney B, Schell M (1990) Inactivation of multiple virulence genes reduces the ability of *Pseudomonas solanacearum* to cause wilt symptoms. Mol Plant-Microbe Interact 3: 293-300.
  27. Liu H, Zhang S, Schell MA, Denny TP (2005) Pyramiding unmarked deletions in *Ralstonia solanacearum* shows that secreted proteins in addition to plant cell-wall-degrading enzymes contribute to virulence. Mol Plant-Microbe Interact 18: 1296-1305.
  28. Denny T, Baek S (1991) Generic evidence that extracellular polysaccharide is a virulence factor of *Pseudomonas solanacearum*. Mol Plant-Microbe Interact 4: 198-206.
  29. Kao CC, Barlow E, Sequeira L (1992) Extracellular polysaccharide is required for wild-type virulence of *Pseudomonas solanacearum*. J Bacteriol 174: 1068-1071.
  30. Saile E, McGarvey JA, Schell MA, Denny TP (1997) Role of Extracellular Polysaccharide and Endoglucanase in Root Invasion and Colonization of Tomato Plants by *Ralstonia solanacearum*. Phytopathology 87: 1264-1271.
  31. Vasse J, Genin S, Frey P, Boucher C, Brito B (2000) The hrpB and hrpG regulatory genes of *Ralstonia solanacearum* are required for different stages of the tomato root infection process. Mol Plant Microbe Interact 13: 259-267.
  32. Kanda A, Yasukohchi M, Ohnishi K, Kiba A, Okuno T, et al. (2003) Ectopic expression of *Ralstonia solanacearum* effector protein PopA early in invasion results in loss of virulence. Mol Plant Microbe Interact 16: 447-455.
  33. Zolobowska L, Van Gijsegem F (2006) Induction of lateral root structure formation on petunia roots: A novel effect of GMI1000 *Ralstonia solanacearum* infection impaired in Hrp mutants. Mol Plant Microbe Interact 19: 597-606.
  34. Meng F, Yao J, Allen C (2011) A MotN mutant of *Ralstonia solanacearum* is hypermotile and has reduced virulence. J Bacteriol 193: 2477-2486.
  35. Schell MA (2000) Control of virulence and pathogenicity genes of *Ralstonia solanacearum* by an elaborate sensory network.Annu Rev Phytopathol 38: 263-292.
  36. Brumbley SM, Carney BF, Denny TP (1993) Phenotype conversion in *Pseudomonas solanacearum* due to spontaneous inactivation of PhcA, a putative LysR transcriptional regulator. J Bacteriol 175: 5477-5487.
  37. Bhatt G, Denny TP (2004) *Ralstonia solanacearum* iron scavenging by the siderophorestaphyloferrin B is controlled by PhcA, the global virulence regulator. J Bacteriol 186: 7896-7904.
  38. Brumbley SM, Denny TP (1990) Cloning of wild-type *Pseudomonas solanacearum* phcA, a gene that when mutated alters expression of multiple traits that contribute to virulence. J Bacteriol 172: 5677-5685.
  39. Clough SJ, Flavier AB, Schell MA, Denny TP (1997) Differential Expression of Virulence Genes and Motility in *Ralstonia (Pseudomonas) solanacearum* during Exponential Growth. Appl Environ Microbiol 63: 844-850.
  40. Flavier AB, Ganova-Raeva, L.M., Schell, M.A., and Denny, T.P. (1997) Hierarchical autoinduction in *Ralstonia solanacearum*: control of acyl-homoserine lactone production by a novel autoregulatory system responsive to 3-hydroxypalmitic acid methyl ester. J Bacteriol 179: 7089-7097.
  41. Genin S, Brito B, Denny TP, Boucher C (2005) Control of the *Ralstonia solanacearum* Type III secretion system (Hrp) genes by the global virulence regulator PhcA. FEBS Lett 579: 2077-2081.
  42. Huang J, Carney BF, Denny TP, Weissinger AK, Schell MA (1995) A complex network regulates expression of eps and other virulence genes of *Pseudomonas solanacearum*. J Bacteriol 177: 1259-1267.
  43. Flavier AB, Clough SJ, Schell MA, Denny TP (1997) Identification of 3-hydroxypalmitic acid methyl ester as a novel autoregulator controlling virulence in *Ralstonia solanacearum*.. MolMicrobiol 26: 251-259.
  44. Flavier AB, Schell MA, Denny TP (1998) An RpoS (sigmaS) homologue regulates acylhomoserine lactone-dependent autoinduction in *Ralstonia solanacearum*. Mol Microbiol 28: 475-486.
  45. Genin S, Boucher C (2004) Lessons learned from the genome analysis of *Ralstonia solanacearum*. Annu Rev Phytopathol 42: 107-134.
  46. Aldon D, Brito B, Boucher C, Genin S (2000) A bacterial sensor of plant cell contact controls the transcriptional induction of *Ralstonia solanacearum* pathogenicity genes. EMBO J 19: 2304-2314.
  47. Marena M, Brito B, Callard D, Genin S, Barberis P, et al. (1998) PrhA controls a novel regulatory pathway required for the specific induction of *Ralstonia solanacearum* hrp genes in the presence of plant cells. MolMicrobiol 27: 437-453.
  48. Brito B, Aldon D, Barberis P, Boucher C, Genin S (2002) A signal transfer system

- through three compartments transduces the plant cell contact-dependent signal controlling *Ralstonia solanacearum* hrp genes. Mol Plant-Microbe Interact 15: 109-119.
49. Brito B, Marena M, Barberis P, Boucher C, Genin S (1999) prhJ and hrpG, two new components of the plant signal-dependent regulatory cascade controlled by PrhA in *Ralstonia solanacearum*. MolMicrobiol 31: 237-251.
50. Valls M, Genin S, Boucher C (2006) Integrated regulation of the type III secretion system and other virulence determinants in *Ralstonia solanacearum*. PLoS Pathog 2: e82.
51. Occhialini A, Cunnac S, Reymond N, Genin S, Boucher C (2005) Genome-wide analysis of gene expression in *Ralstonia solanacearum* reveals that the hrpB gene acts as a regulatory switch controlling multiple virulence pathways. Mol Plant Microbe Interact 18: 938-949.
52. Mukaihara T, Tamura N, Murata Y, Iwabuchi M (2004) Genetic screening of Hrp type III-related pathogenicity genes controlled by the HrpB transcriptional activator in *Ralstonia solanacearum*. Mol Microbiol 54: 863-875.
53. Cunnac S, Boucher C, Genin S (2004) Characterization of the cis-acting regulatory element controlling HrpB-mediated activation of the type III secretion system and effector genes in *Ralstonia solanacearum*. J Bacteriol 186: 2309-2318.